



# Structural preferences and multiple bonding interactions in hetero-*closo*-dodecaborates of group-14 elements – A theoretical study

Brindha Veerappan, Krishnamoorthy Bellie Sundaram  

Department of Chemistry (SF), PSG College of Arts and Science, Coimbatore, Tamil Nadu, 641014, India

Received 21 August 2025, Revised 29 October 2025, Accepted 3 November 2025, Available online 5 November 2025, Version of Record 9 November 2025.

 [What do these dates mean?](#)



Show less 

 Outline |  Share  Cite

<https://doi.org/10.1016/j.jmkgm.2025.109214> 

[Get rights and content](#) 

Full text access


## Highlights

- DFT computations on the dimers of group 14 hetero-*closo*-dodecaborates resulted in stable clusters except carbon.
- An intra-cage electron delocalization improves the stability of the dimers.
- The dimers of tin clusters with 'Sn-B' bond has 1.3eV more stable than that of 'Sn=Sn' bond.
- An ELF analysis shows has strong bonding with good electron density for the cluster (2).
- The predominant dimer formation with 'Pb=Pb' multiple bond promises their application in solid state electrolyte.

## Abstract

The geometric and electronic structural features of group-14 hetero-*closo*-dodecaborates such as monomers  $[\text{HEB}_{11}\text{H}_{11}]^-$ ,  $[\text{EB}_{11}\text{H}_{11}]^{2-}$  [ $\text{E}=\text{C}$  (**1a**), Si (**2a**), Ge (**3a**), Sn (**4a**) and Pb (**5a**)] and dimers  $[(\text{EB}_{11}\text{H}_{11})_2]^{2-}$  [ $\text{E}=\text{C}$  (**1**), Si (**2**), Ge (**3**), Sn (**4**) and Pb (**5**)] analyzed using density functional theory (DFT) method at BP86/Def2-TZVP level of theory. The structural features such as the geometry, metrical parameters and bonding interactions are theoretically studied. The dispersion correction method has been used to get more insight which improves the metrical parameter of group-14 elements towards experimental parameters. The electronic structure of the clusters has been studied using frontier molecular orbital analysis (FMO). It shows the group-14 elements electronic contribution and possible electron delocalization between the cages to stabilize the dimeric clusters. DFT (BP86/Def2-TZVP) calculations on the clusters confirm the thermal stability of the newly modelled dimeric clusters (**2–5**) which could have diverse applications. The  $^{13}\text{C}$ ,  $^1\text{H}$ ,  $^{29}\text{Si}$ , and  $^{11}\text{B}$  nmr chemical shift values computed at DFT (BP86/Def2-TZVP) level are highly useful in identifying and assigning the individual atoms of the monomeric and dimeric clusters. From the global reactivity descriptors and local descriptors analysis, an identification of reactive sites with respect to electrophilic and nucleophilic centers of the reactant has been reported.

## Graphical abstract



[Download: Download high-res image \(291KB\)](#)

[Download: Download full-size image](#)



Previous

Next



## Keywords

$[\text{EB}_{11}\text{H}_{11}]^{2-}$ ;  $[(\text{EB}_{11}\text{H}_{11})_2]^{2-}$ ; DFT-D3; Group-14; Stanna-*closo*-dodecaborate

## 1. Introduction

Multiple bonding interactions between group 14 elements have a unique electronic structure and bonding [[1], [2], [3], [4], [5]] which attracts most of the theoretical researchers especially through the density functional theory (DFT) method [[6], [7], [8]]. The bonding interactions of RE/ER [ $(/ \rightarrow -, =, \equiv)$ , ( $\text{E}=\text{C}$ , Si, Ge, Sn, Pb)] molecules are studied by Shigeru Nagase et al. [9] The heavier group multiple bonds are calculated by using Grimme's dispersion correction D3 method [10]. Icosahedral heteroborates are the kind of Lewis acid-base in which the electrophilicity or nucleophilicity of the clusters mainly depends upon the heteroatoms [11,12]. Boron is a special kind of element that is capable of forming extended binary hydrides, called boranes. This precise species was discovered by Alfred Stock in 1912 [13]. In contrast to hydrocarbons, boranes avoid the formation of chain structures and clearly prefer the formation of polyhedral clusters to overcome the electron deficiency. It can form 2c-2e, 3c-2e, and 4c-2e bonding. This architecture of boranes was proposed by Lipscomb in 1963, and the Nobel Prize was awarded to him for his excellent work on explaining the bonding in these boranes in 1976 [[14], [15], [16]]. Icosahedral borates  $[\text{B}_{12}\text{H}_{12}]^{2-}$  are the twelve vertices of BH units exhibiting perfect  $A_g$  symmetry, these dianionic *closo*-dodecaborates

arise from the symmetric arrangement of electrons throughout the molecular orbitals of the 12 vertices. Icosahedral borate structures are a kind of three-dimensional configurations made up of triangular B-B-B units. The clusters in this category possess different bonding characteristics compared to typical boron cluster compounds. These clusters exhibit sigma aromaticity or three-dimensional aromaticity due to the spatial delocalization of electron density. The delocalization of two surplus electrons in a dianionic center contributes stability to the sigma bonds of the entire cage, resulting in distinctive steric and electronic properties [[17], [18], [19]]. The heteroatom present in the apical vertex causes slight distortion in  $I_h$  symmetry. These distorted icosahedral hetero-*closo*-dodecaborate clusters possess very unique structural characteristics. The Todd group found the first organometallic derivative of dianionic dodecaborate in 1992 [20]. The lone pair electrons at the tin vertex impart Lewis acidity to the stanna-*closo*-dodecaborates, resulting in reactivity towards Lewis bases. It functions as a strong sigma donor and a notable  $\pi$  acceptor, which contributes to ligation, thermodynamic trans influence, and so forth [17,18]. These icosahedral clusters have been used in boron neutron capture therapy (BNCT), HIV inhibitors, solid state electrolytes, super acids, ligation and etc. [[21], [22], [23], [24], [25], [26], [27], [28], [29], [30], [31], [32], [33]].

Hetero-*closo*-dodecaborate (C, Si, Ge, Sn and Pb) can possess outstanding superionic conductivity, especially above the transition temperatures at which they occur. These anionic *closo*-dodecaborates are found in salts like  $\text{LiCB}_{11}\text{H}_{12}$ ,  $\text{NaCB}_{11}\text{H}_{12}$  and  $\text{Li}_2\text{PbB}_{11}\text{H}_{11}$  which have excellent ionic conductivity and frequently surpass that of liquid electrolytes. At high temperatures, the dodecaborate structure of these salts changes, causing phase transitions that improve superionic conductivity. By removing the possibility of leaks and combustible components, solid-state electrolytes provide a safer substitute for liquid electrolytes. Hetero-*closo*-dodecaborate salts enhance ionic conductivity may contribute to batteries increased energy density. Due to their strong ionic conductivity, better cycle performance, long life and possibility for property tuning, hetero-*closo*-dodecaborates show promise as a novel class of solid-state electrolytes for rechargeable batteries. The goal of ongoing research is to optimize them for real-world use in next-generation batteries and increase their stability, especially at room temperature [[26], [27], [28], [29]].


Boron drugs are specifically designed to target the tumor cells of the patient. The drugs are administered via intravenously, the beam of slow neutrons passes. Boron atoms have the better cross section to capture the neutron particles, then dissociate into lithium ions and high energy alpha particles. It has high linear energy transfer (LET), which is a measure of biological effectiveness of radiation in a more localized manner. So, it can specifically kill the tumor cells, with the minimum damage of healthy cells present in the adjacent areas. Currently, Boron delivery agent researchers are fueling continuous creativity towards better activity and clinical trials with minimum damage [[23], [24], [25], [26]].

The local descriptor and global reactive descriptors are key factors to determine the chemical nature of the system [[34], [35], [36]]. The amphiphilic nature is that the system has both nucleophilic and electrophilic nature which can also be calculated through computational methods [[37], [38], [39]]. This factor is mainly determined by Fukui functions, the global reactive descriptors such as chemical potential ( $\mu$ ), hardness ( $\eta$ ), softness ( $S$ ) and electrophilicity ( $\omega$ ). These descriptors can be calculated by using DFT method [38,40].

The chemical softness and hardness play a major role in the behavior of the system, this concept first introduced by Robert S. Mulliken at 1952 [41]. Then Hard and Soft Acid Base (HSAB) principle evolved by Pearson postulates. The absolute hardness was introduced by Pearson and Parr. Then Pearson published maximum hardness principle MHP, these two MHP and HSAB are used for the hardness and chemical potential. The Koopmann's theorem mainly used to derive the global reactive descriptors [42]. According to Koopmann's theorem, we can calculate the adiabatic ionization potential (AIP) and vertical ionization potential (VIP). The first ionization energy is derived from the highly occupied molecular orbitals (HOMO), the negative value of the highly occupied molecular orbital's energy plus the electrostatic potential energy of the electron removed from the system. Similarly, we can calculate the ionization potential (IP) average of the highly occupied molecular orbitals (HOMO) of the neutral system (N) and the lowest unoccupied molecular orbitals (LUMO) of the cationic system (N-1) [43].

A dimeric cluster of group-14 icosahedral geometry is reported in 2015 by Wesemann et al., The first synthesized hetero-*closo*-dodecaborate dimer is  $[\text{Bu}_3\text{MeN}]_2 [(\text{GeB}_{11}\text{H}_{11})_2]$ , a colourless crystalline solid synthesized by the

oxidation of germa-*closo*-dodecaborate using iodine and ferroceniumhexafluorophosphate. The dimer is dianionic in nature with the structural formula  $[(\text{GeB}_{11}\text{H}_{11})_2]^{2-}$  [44]. Here we have studied the monomers **1a-5a** and their corresponding dimers **1-5** using the DFT method to analyze their molecular structural features, stability and bonding interactions (Scheme 1). Computational chemistry techniques are being used as valuable tools to help with both the modeling of novel compounds and to study the complete structural characterization of existing compounds [[45], [46], [47], [48], [49], [50]]. The following approaches were used to study each of the clusters: the geometry optimization by DFT method at BP86/Def2-TZVP level, molecular orbital analysis, frequency computation, nmr property calculation and reactivity descriptor analysis.



Download: [Download high-res image \(348KB\)](#)

Download: [Download full-size image](#)

Scheme 1. The schematic diagram representing the molecules studied [E=C (1a), Si (2a), Ge (3a), Sn (4a), Pb (5a)].


## 2. Computational details

The hetero-*closo*-dodecaborates are optimized by using BP86/Def2-TZVP level and the interactions are studied by D3BJ method. All of the density functional theory computations in this work were done using ORCA software developed by F. Neese and colleagues [51]. From the Vosko-Wilk-Nusair parameterization [52], the Becke 88 gradient correction for exchange and Perdew 86 correlation inside the local density approximation (LDA) were applied [[53], [54], [55]]. The DFT functional BP86 highly favors the theoretical values with experimental values in metallaboranes [12,[56], [57], [58]]. For each clusters, the Def2-TZVP (triple zeta valence with polarization function) basis set with ZORA (zeroth order regular approximation) was employed. TightSCF convergence requirements were used for all calculation [59]. The bonding interactions of group-14 elements are analyzed by using DFT-D3 method. The Grimme's dispersion correction D3 [60] with BJ (Beck-Johnson) damping [[61], [62], [63]] has been used to approach the closer value of experimental values for the heavy atom system [9,10]. An electron localization function (ELF) analysis done by using ORCA and Multiwfn analyzer. Using the ORCA software's EPRNMR module, the DFT-optimized geometries were utilized to compute nmr parameters such as chemical shifts and shielding constants [64]. The tetramethylsilane (TMS) has been used as reference compound, for computing  $^1\text{H}$  and  $^{13}\text{C}$  nmr chemical shift values [65]. The calculated  $^{11}\text{B}$  nmr chemical shift values were translated to the standard  $\text{BF}_3 \cdot \text{OEt}_2$  scale using the experimental value of +16.6ppm for  $\text{B}_2\text{H}_6$ , with  $\text{B}_2\text{H}_6$  serving as the major reference point [65,66]. The idea of generalized philicity was presented by Chattaraj et al. It includes information on the electrophilic/nucleophilic power of a specific atomic location in a molecule, as well as nearly all of the previously known various global and local reactivity and selectivity characteristics [67,68]. Further, local descriptors on cluster systems are also explored through DFT method. The binding site of atomic hydrogen on silicon clusters using the local descriptor Fukui Functions [69] and they found the criteria of "maximum matching" between the Fukui functions to predict the best interaction between small ( $\text{Si}_2$ - $\text{Si}_6$ ) silicon clusters to form larger ( $\text{Si}_4$ - $\text{Si}_8$ ) ones [70]. DFT methods have already been used to find out stable clusters and molecules in the potential energy surface, by kick-Fukui method [71]. The stable global minimum structures of  $\text{Na}(\text{Si}_{5-n}(\text{BH})_n)^-$  and  $\text{Si}_{5-n}(\text{BH})_n^{2-}$  ( $n=0-5$ ) cluster systems are explored successfully using DFT method by substituting  $\text{NaSi}_5^-$  and  $\text{Si}_5^{2-}$  for B-H units [72].

## 3. Results and discussion

### 3.1. Geometrical structure

The geometrical structures of the group-14 clusters are studied by using BP86/Def2-TZVP level. The optimized geometries are given in Fig. 1. We have optimized monomers and its respective dimers of the group-14 *closo*-dodecaborates. The DFT (BP86/Def2-TZVP) computed metrical parameters and dispersion corrected values of heteroborate monomers (**1a-5a**) are provided in Table S1 and Table S2 respectively. Similarly the DFT (BP86/Def2-TZVP) computed metrical parameters and dispersion corrected values of heteroborate dimers (**1-5**) are provided in Table S3 and Table S4 respectively. The bonding distance of E (C, Si, Ge, Sn and Pb) and B atoms are around 1.7091 Å for [HCB<sub>11</sub>H<sub>11</sub>]<sup>-</sup> (**1a**), 2.1346 Å for [SiB<sub>11</sub>H<sub>11</sub>]<sup>2-</sup> (**2a**), 2.0326 Å [2.0241 Å] for [MeSiB<sub>11</sub>H<sub>11</sub>]<sup>1-</sup> (**2b**) [73], 2.2204 Å for [GeB<sub>11</sub>H<sub>11</sub>]<sup>2-</sup> (**3a**), 2.4029 Å [2.389 (3) Å] [74] for [SnB<sub>11</sub>H<sub>11</sub>]<sup>2-</sup> (**4a**) and 2.4917 Å for [PbB<sub>11</sub>H<sub>11</sub>]<sup>2-</sup> (**5a**). The increasing order of bond length is **1a** < **2a** < **3a** < **4a** < **5a** < **2b**, the increasing order depends upon the increasing the radii of group-14 elements and interatomic repulsions between B and group-14. The increasing order of bond interactions between E and B atoms using D3BJ method is 1.7061 Å (**1a**) < 2.1307 Å (**2a**) < 2.2143 Å (**3a**) < 2.3973 Å [2.389 (3) Å] (**4a**) < 2.4822 Å (**5a**) < 2.0276 Å [2.0241 Å] (**2b**). Both DFT-D3 calculated and x-ray crystal structure E-B bond distance of [MeSiB<sub>11</sub>H<sub>11</sub>]<sup>1-</sup> (**2b**) is 2.02–2.03 Å and [SnB<sub>11</sub>H<sub>11</sub>]<sup>2-</sup> (**4a**) is 2.38–2.40 Å. The DFT-D3 calculated cage vertical distances [CVD] are 3.2286 (**1a**) < 3.9071 (**2a**) < 4.0052 (**3a**) < 4.2298 (**4a**) < 4.3346 (**5a**), the CVD values are gradually increasing from carborate to plumbaborate cage. The dispersion correction method (D3BJ) improves the bonding distances, which are closer to experimental values. The bond angles of B1-E-B2 are 62.834° (**1a**) > 54.904 [54.488]° (**2b**) > 50.578° (**2a**) > 48.974° (**3a**) > 45.502 [45.380]° (**4a**) > 43.998° (**5a**). The DFT-D3 method B1-E-B2 bond angles are 62.849 (**1a**) > 54.934 [54.488] (**2b**) > 50.576 (**2a**) > 48.976 (**3a**) > 45.522 [45.380] (**4a**) > 44.043 (**5a**). The cluster (**2b**) and (**4a**) have a good agreement with an experimental value [73,74].




[Download: Download high-res image \(1MB\)](#)

[Download: Download full-size image](#)

Fig. 1. DFT optimized geometries of group-14 hetero-*closo*-dodecaborate monomers (**1a – 5a**) and dimers (**1–5**) at BP86/Def2-TZVP level.

The DFT computed dimeric clusters are  $[(CB_{11}H_{11})_2]^{2-}$ (**1**),  $[(SiB_{11}H_{11})_2]^{2-}$ (**2**),  $[(GeB_{11}H_{11})_2]^{2-}$ (**3**),  $[(SnB_{11}H_{11})_2]^{2-}$ (**4A**),  $[(SnB_{11}H_{10})(SnB_{11}H_{11})_2]^{3-}$ (**4B**) and  $[(PbB_{11}H_{11})_2]^{2-}$ (**5**) have E/E bonding interactions with 1.5535, 2.3290, 2.4048, 2.7377 and 2.8687 Å for the clusters (**1–5**) respectively. The dispersion corrected E/E interactions are 1.5386, 2.2955, 2.3715, 2.6986 and 2.8236 Å. An experimental values of the E/E bond of Germa-*closo*-dodecaborate dimer  $[Bu_3MeN]_2 [(GeB_{11}H_{11})_2]$  is 2.329 (1) Å, which is very closer to the DFT-D3 computed value of  $[(GeB_{11}H_{11})_2]^{2-}$  (**3**) is 2.3715 Å, the remaining compounds are not synthesized [44]. So, we have compared an experimental value of the related compounds for the clusters (**1**), (**2**), (**4A**), (**4B**) and (**5**). The C-C interaction of cluster (**1**) is 1.5386 Å which similar to the alkanes C-C bond distance around 1.540 Å [75], confirming the single bond nature. DFT computed

Si=Si bond length of 2.2955 Å is similar to the various disilene distance 2.304 (2) Å of 1-methyl-2-phenyl-1,2-disilacloso-dodecaborane (12) [(PhSi)(MeSi)B<sub>10</sub>H<sub>10</sub>] [76], the distance 2.2294 (11) of [L<sub>2</sub>Si=SiL<sub>2</sub>] (L=:C [N (2,6-Pr<sup>i</sup><sub>2</sub>C<sub>6</sub>H<sub>3</sub>)CH]<sub>2</sub>) [77], 216 p.m. or 2.1600 Å presents in (Mes)<sub>2</sub>Si=Si(Mes)<sub>2</sub> [Mes – mesityl group (C<sub>6</sub>H<sub>2</sub>-2,4,6-Me<sub>3</sub>)] [78,79]. The DFT-D3 and DFT computed Sn=Sn double bond distance 2.6986 and 2.7377 Å is very close to the stannylene or distanene (Sn=Sn) double bond 2.6683 (10) Å presents in (<sup>t</sup>Bu<sub>2</sub>MeSi)<sub>2</sub>Sn=Sn(SiMe<sup>t</sup>Bu<sub>2</sub>)<sub>2</sub> [[80], [81]], and 2.702 in ((Me<sub>3</sub>Si)<sub>3</sub>Si)<sub>2</sub>Sn=Sn(2,4,6-Me<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>)<sub>2</sub> [84]. The experimentally available structure for stannaborate is [(SnB<sub>11</sub>H<sub>11</sub>)(SnB<sub>11</sub>H<sub>10</sub>)]<sup>3-</sup> (**4B**) [44], shown in Fig. 2 and the two stannaborate clusters are inter-connected by 'Sn-B' bonding with 2.2877 Å inter atomic distance. The dispersion corrected (D3BJ) 'Sn-B' bond distance is 2.2445 Å. Both of these DFT and DFT-D3 Sn-B distance is smaller than the usual Sn-B bonds in our monomers and experimental structures [12,74]. The DFT-D3 and DFT computed Pb=Pb bond length of 2.8236 Å and 2.8687 Å is similar to the Pb=Pb bond length 2.833 Å of L:Pb=Pb:L, 2.844 Å of Ph<sub>3</sub>Pb-PbPh<sub>3</sub> and 2.903 Å of Si(SiMe<sub>3</sub>)<sub>3</sub>MesPb=PbSi(SiMe<sub>3</sub>)<sub>3</sub>Mes (Mes=2, 4, 6 - trimethylphenyl) [9].



Download: [Download high-res image \(610KB\)](#)

Download: [Download full-size image](#)

Fig. 2. DFT optimized geometry of [(SnB<sub>11</sub>H<sub>11</sub>)(SnB<sub>11</sub>H<sub>10</sub>)]<sup>3-</sup> (**4B**) cluster at BP86/Def2-TZVP level.

The B1-E1-E1 angles are gradually increasing from the left to right of the C to Pb series, 119.485° (**1**) < 128.660° (**2**) < 130.149° [128.095°] (**3**) < 134.409° (**4A**) < 137.629° (**5**). The cluster (**3**) has closer value with experimental values. The E1-B1-E2 bond angle of cluster (**4B**) is 177.466°. All the DFT and dispersion corrected DFT-D3 binding interactions, bond length and bond angles are in good agreement with experimental values and related experimental values (Table S3 and S4).

## 3.2. Chemical descriptors

The chemical descriptors are the indicators of the molecule nature such as reactivity and reactive center. The frontier molecular orbitals (FMO) of the clusters studied are provided in Figure SF3-Figure SF5. Those are divided into two categories, local descriptors like highly occupied molecular orbitals (HOMO), lowest unoccupied molecular orbitals (LUMO), energy gap ( $\Delta E$ ) and global reactive descriptors like chemical potential ( $\mu$ ), hardness ( $\eta$ ), softness ( $S$ ), electrophilicity ( $\omega$ ), dipole moment, ionization potential ( $I_p$ ), electron affinity ( $E_A$ ). These parameters are calculated by using  $E_{\text{HOMO}}$  and  $E_{\text{LUMO}}$ . All these descriptors are the major key factor for identifying the natures of the clusters. These descriptors are derived by using Koopmann's theorem.

### 3.2.1. Local descriptors

We have computed (Table 1) the descriptors for monomers (**1a**, **2a**, **2b**, **3a**, **4a** and **5a**) and dimers (**1**, **2**, **3**, **4A**, **4B** and **5**). The energy gap determines the reactivity and kinetic stability of the geometry, energy of the group-14 monomers are around 3.8368–6.4526 eV. The energy gap range of group-14 dimer is 1.5361–6.3217 eV with highest  $E_{\text{LUMO-HOMO}}$  energy gap values of 6.3217 eV for carborate dimer cluster **1**. The increasing order of energy gap for monomer is **5a** < **4a** < **3a** < **2a** < **1a** and the dimer is **5** < **4A** < **4B** < **3** < **2** < **1**. The increasing order shows that the size of the group-14 element increases the energy gap of the clusters decreases.

Table 1. DFT calculated chemical descriptors (eV) of the monomers (**1a-5a**) and dimeric clusters (**1-5**) at BP86/Def2-TZVP level.

| Cluster                      | 1a      | 1       | 2a      | 2       | 3a      | 3       | 4a      | 4A      | 4B      | 5a      | 5       |
|------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| HOMO (eV)                    | -3.3197 | -0.9054 | 2.3009  | -1.4080 | 1.9679  | -1.2635 | 1.7134  | -0.9958 | 2.7506  | 1.9674  | -0.7430 |
| LUMO (eV)                    | 3.1329  | 5.4163  | 6.7607  | 3.6702  | 6.8277  | 3.0466  | 5.9550  | 1.8011  | 6.8949  | 5.8040  | 0.7931  |
| E <sub>LUMO - HOMO</sub>     | 6.4526  | 6.3217  | 4.4598  | 5.0782  | 4.8598  | 4.3101  | 4.2416  | 2.7969  | 4.1443  | 3.8368  | 1.5361  |
| Chemical potential( $\mu$ )  | -0.0934 | 2.2555  | 4.5308  | 2.2622  | 4.3978  | 1.8116  | 3.8342  | 0.4026  | 4.8228  | 3.8858  | 0.0251  |
| Hardness( $\eta$ )           | 3.2263  | 3.1609  | 2.2299  | 2.5391  | 2.4299  | 2.1551  | 2.1208  | 1.3985  | 2.0722  | 1.9184  | 0.7681  |
| Softness (S)                 | 0.3100  | 0.3164  | 0.4485  | 0.3938  | 0.4115  | 0.4640  | 0.4715  | 0.7151  | 0.4826  | 0.5213  | 1.3019  |
| Electrophilicity( $\omega$ ) | 0.0014  | 0.8047  | 4.6029  | 1.0077  | 3.9797  | 0.7614  | 3.4659  | 0.0580  | 5.6122  | 3.9354  | 0.0004  |
| Dipole Moment (Debye)        | 2.65    | 0.00    | 0.42    | 0.03    | 5.49    | 0.07    | 9.14    | 0.11    | 8.38    | 14.11   | 0.17    |
| Ionization potential(eV)     | 3.3197  | 0.9054  | -2.3009 | 1.4080  | -1.9679 | 1.2635  | -1.7134 | 0.9958  | -2.7506 | -1.9674 | 0.7430  |
| Electron Affinity (eV)       | -3.1329 | -5.4163 | -6.7607 | -3.6702 | -6.8277 | -3.0466 | -5.9550 | -1.8011 | -6.8949 | -5.8042 | -0.7931 |

Fukui function has been used to calculate the local descriptors and reactivity sites. Hirsh field analysis lead to the local philicity of the system from which we can determine either it is nucleophilic, electrophilic, neutral or amphiphilic in nature. The following equations 1, 2, 3 have been used to calculate nucleophilic, electrophilic and radical attack [38].

$$f_k^+ = q_k(N+1) - q_k(N) \quad 1$$

$$f_k^- = q_k(N) - q_k(N-1) \quad 2$$

$$f_k^0 = [q_k(N+1) - q_k(N-1)]/2 \quad 3$$

The dimerization of germa-*closo*-dodecaborate gives the dimeric 'Ge=Ge' bonded [(GeB<sub>11</sub>H<sub>11</sub>)]<sup>2-</sup> (**3**) cluster. But, in the case of stanna-*closo*-dodecaborate produces the 'Sn-B' bonded [(SnB<sub>11</sub>H<sub>11</sub>)(SnB<sub>11</sub>H<sub>10</sub>)]<sup>3-</sup> (**4B**) cluster. The quantum chemical computation favors the remaining group-14 dimeric product. The stanna-*closo*-dodecaborate computation possess the favorable results for both the 'Sn=Sn' (**4A**) and 'Sn-B' (**4B**) bonded cluster system. But, the energy gap for the 'Sn-B' system is 4.13eV with 8.55 Debye dipole moment and 2.7920eV for the 'Sn=Sn' bonded system. The energy difference for these two systems is around 1.3eV. So, we have done Fukui function analysis to confirm the product formation.

The calculation of electrophilic Fukui function ( $f_k^+$ ), nucleophilic Fukui function ( $f_k^-$ ) and neutral Fukui function ( $f_k^0$ ) through N-1, N+1 and N systems. The calculation shows that the  $f_k^-$ ,  $f_k^0$  favours the B2 vertex with -0.0190, -0.0197 and  $f_k^+$  favours B4 vertex with -0.0197. So, B2 and B4 vertex is electrophilic and nucleophilic respectively, in nature. So, this analysis highly favours the 'Sn-B' bonded [(SnB<sub>11</sub>H<sub>11</sub>)(SnB<sub>11</sub>H<sub>10</sub>)]<sup>3-</sup> (**4B**) cluster formation. But, in the tin vertex, the calculations gives  $f_k^+$ ,  $f_k^-$  and  $f_k^0$  -0.5402, -0.2689 and -0.4046. Hence, tin vertex is highly nucleophilic in nature and less electrophilic nature. So, the 'Sn=Sn' (**4A**) bonded dimeric cluster is less favored.

### 3.2.2. Global reactive descriptors

The global reactive descriptors are calculated by using below equations (4)-(8), the equation solved by using frontier molecular orbital HOMO and LUMO [45]. The DFT computed chemical descriptors are given in Table 1.

$$\text{Electronegativity} = -\frac{1}{2} (E_{\text{LUMO}} + E_{\text{HOMO}}) \quad 4$$

$$\text{Global Chemical Potential } (\mu) = \frac{1}{2} (E_{\text{LUMO}} + E_{\text{HOMO}}) \quad 5$$

$$\text{Global Hardness } (\eta) = \frac{1}{2} (E_{\text{LUMO}} - E_{\text{HOMO}}) \quad 6$$

$$\text{Global Softness } (S) = 1/\eta \quad 7$$

$$\text{Global Electrophilicity } (\omega) = \frac{\mu^2}{2\eta} \quad 8$$

Chemical potential ( $\mu$ ) of the carboratecluster (**1a**) and plumbaborate (**5**) has very lower values  $-0.0934$  and  $0.0251$ , the lowest value obtained due to higher nucleophilicity of the cluster. The monomers have the higher ( $\mu$ ) value than the dimers except carborate system. It determines that the dimers are highly nucleophilic in nature than the monomers; the heavy group clusters Sn and Pb dimers are good at this criterion.

Hardness ( $\eta$ ) determines the overall stability of the clusters. The hardness increases the stability of the system increases. The hardness of the carborates (**1, 1a**) has higher value around  $3.2263$  to  $3.1609$  than the remaining group-14 clusters. But, in the case of electrophilicity ( $\omega$ ), the carborates (**1, 1a**) have lower value than the remaining group-14 clusters. The hardness values are gradually decreases from the silaborates to plumbaborates except (**4B**).

Softness ( $S$ ) of the clusters is lower in all geometries except the dimeric plumbaborate (**5**) geometry as  $1.3019\text{eV}$ . The higher softness value indicates that the cluster has the possibility to better reactivity. Similarly, the better reactivity of this cluster is supported by the lower energy gap than the remaining group-14 geometries with less hardness and electrophilicity. Hence, we can find the stability and reactivity of the clusters through hardness ( $\eta$ ) and softness ( $S$ ).

Electrophilicity( $\omega$ ) of the monomeric clusters (**2a, 2, 3a, 4a, 4B** and **5a**) are higher than the dimers except carborates. In the dimeric system, the silaborate only has the considerable electrophilicity value at  $1.0077\text{eV}$ . The dipole moment of the dimeric clusters (**1, 2, 3, 4** and **5**) are almost zero except dimer (**4B**) with  $8.3763$  Debye dipole moment and the monomers are in the order of  $14.1067$  (**5a**) $>9.1407$  (**4a**) $>5.4925$  (**3a**) $>2.6467$  (**1a**) $>0.4212$  (**2a**). Among the monomers studied the plumba-*closo*-dodecaborate (**5a**) polar cluster has highest dipole moment (DM) value of  $14.11$  Debye (D) when compared to the than the remaining group-14 monomeric clusters. According to Koopmann's theorem negative value of HOMO is termed as ionization potential (IP) and negative value of LUMO is termed as electron affinity (EA). The carborate shows the highest IP  $3.3197\text{eV}$  and dimeric plumbaborate shows highest EA  $-0.7931\text{eV}$ .

### 3.3. Polyhedral skeletal electron pair theory (PSEPT) and cluster valence electron (CVE) count

The skeletal electron pair count and electronic structure of the clusters are studied by PSEPT theory [[82], [83], [84]]. The PSEPT and CVE suggests that the icosahedral borate  $[\text{B}_{12}\text{H}_{12}]^{2-}$  cluster has 13 skeletal electron pair (SEP) possess the *closo* geometry, with the general formula of the *closo* geometry  $[\text{B}_n\text{H}_n]^{2-}$  [82].

$$12 \times 2 (\text{BH}) + 2 (2\text{e}^-) = 26 \text{ e}^- \text{ s}$$

$$26/2 = 13$$

$$\text{SEP} = (n + 1) = (12 + 1) = 13$$

$$n = 12 \text{ (12 - vertex icosahedral geometry)}$$

Similarly, the group 14 clusters also possess the *closo* geometry with 13 SEP and  $n=12$ . So, the monomers of the group 14 clusters obeys the wades rule, the geometrical formula of the group 14 clusters are  $[\text{HCB}_{11}\text{H}_{11}]^-$ ,  $[\text{SiB}_{11}\text{H}_{11}]^{2-}$ ,  $[\text{GeB}_{11}\text{H}_{11}]^{2-}$ ,  $[\text{SnB}_{11}\text{H}_{11}]^{2-}$  and  $[\text{PbB}_{11}\text{H}_{11}]^{2-}$ .

$$11 \times 2 (\text{BH}) + 1 \times 3 (\text{C}) + 1 (1\text{e}^-) = 26 \text{ electrons}$$

$$11 \times 2 (\text{BH}) + 1 \times 2 (\text{E}) + 2 (2\text{e}^-) = 26 \text{ electrons (E = Si, Ge, Sn, Pb)}$$

$$26/2 = 13$$

$$\text{SEP} = (n + 1) = (12 + 1) = 13$$

$$n = 12$$

Hence, the clusters possess *closo* icosahedral geometry, the structure is slightly distorted at the hetero atom vertex. So, the geometrical structure of the group-14 cluster is distorted icosahedral structure and termed as distorted hetero-*closo*-dodecaborate.

### 3.4. Bond order

The Mayer bond analysis used to determine the bond order and charge density of the atoms. The E/E bond orders of dimeric compounds are given in Table 2. The bond order of the E/E bond presents in the carborate (**1**) is 1.1633, 0.8227 in silaborate (**2**), 0.8299 in germaborate (**3**), -3.3836 for stannaborate (**4A**) and 0.8944 for plumbaborate (**5**) clusters. Hence, the bond order of all clusters proves that the bonding interactions presents in the group-14 clusters except stannaborates. The stannaborate shows that the negative bond order value in DFT (-3.3836) and DFT-D3 (-0.3721) method which could be due to the restriction or bugs in the method/software for this specific system. The bond order values obtained from the DFT-D3 methods are slight lower than the DFT method for C and Si systems. But, in the remaining cases, the E-E (Ge, Sn, Pb) bond order is slightly higher than the DFT method. The difference for the Ge cluster is 0.0813 and Sn cluster is 3.0115. The DFT-D3 computed Sn2-B1 bond order of cluster (**4B**) is 0.9361. The Pb cluster is DFT-D3 (1.0439) value is higher than the DFT (0.8944) method. The difference between these two methods proves the D3 method is highly favorable for the heavy atoms.

Table 2. The E/E bond order of group 14 hetero-*closo*-dodecaborate dimers at BP86/Def2-TZVP level using normal DFT and DFT-D3 method.


| Clusters | C    |      | Si   |      | Ge   |      | Sn  |    | Pb   |      |
|----------|------|------|------|------|------|------|-----|----|------|------|
|          | DFT  | D3   | DFT  | D3   | DFT  | D3   | DFT | D3 | DFT  | D3   |
| Mayer    | 1.16 | 1.04 | 0.82 | 0.77 | 0.83 | 0.91 | -   | -  | 0.89 | 1.04 |

### 3.5. Electronic structure

The frontier molecular orbitals (FMO) of the clusters studied are provided in Fig. SF3 - SF5. The highly occupied molecular orbitals (HOMO) of the monomers (**1a-5a**) mainly involve the electron density located at the boron vertices. In contrast, in the lowest unoccupied molecular orbitals (LUMO), electron density is localized at the group 14 elements. In the dimeric clusters, also similar trend continues. In the case of LUMO and LUMO+1 of disilaborate (**2**) electrons are delocalized over the two *closo* clusters, which is seen in the picture Fig. SF3 - SF5. The LUMO+1 and LUMO+2 of germaborates (**3**) has electron density over the Ge-Ge bonding. The LUMO+1 and LUMO+2 of stannaborate (**4A**) and plumbaborate (**5**) have the electron density over the E-E bond. The HOMO of the cluster (**4B**) shows the electron density over the two *closo* clusters, shown in Fig. 2.

### 3.6. Electron localization function (ELF) analysis

Electron Localization Function (ELF) Analysis from computations are highly useful in confirming the metal-metal, multiple bonding interactions [56]. The bonding nature using ELF analysis of group 14 elements presents in the dimeric hetero-*closo*-dodecaborates are shown in Fig. 3. The colour bar range is 0-1, the highest electron localization at red and lowest electron localization at dark blue. The electron cloud value below 0.5 shows the low electron density [56]. The carborate has good electron density between two carbon atoms as red colour range 0.9-1. The order of electron cloud density from carbon to lead is carborate < silaborate > germaborate > stannaborate > plumbaborate. But, in the case of stannaborates electron density, the Sn-B density (0.7-0.8 orange) overcomes Sn-Sn electron density (green 0.6). The overall ELF analysis shows very good electron density of group 14 elements for sila-*closo*-dodecaborate dimer (**2**).



[Download: Download high-res image \(2MB\)](#)

[Download: Download full-size image](#)

Fig. 3. ELF diagram for the DFT optimized geometries of dimers (1–5) at BP86/Def2-TZVP level using Multiwfn analyzer.

### 3.7. Thermochemistry

The thermochemical calculations of the monomers and dimers through frequency analysis give no imaginary frequency and the obtained optimized geometries also minima in the potential energy surface. The DFT (BP86/Def2-TZVP) computed thermal energy, enthalpy, entropy, Gibbs free energy are given in Table 3. The DFT computed free energy of the dimer-forming reactions, confirming the stable nature of the modelled dimers and supports their viable synthesis.

Table 3. DFT calculated thermochemical properties of group 14 hetero-*closo*-dodecaborate monomers and dimers at BP86/Def2-TZVP level.

| Clusters   | Total thermal energy (U) | Enthalpy (H) | Entropy (S) | Gibbs free energy (G) |
|--|--------------------------|--------------|-------------|-----------------------|
| [HCB <sub>11</sub> H <sub>11</sub> ] <sup>-</sup>  | -318.8699Eh              | -318.8689Eh  | 0.0397Eh    | -318.9087Eh           |
| [SiB <sub>11</sub> H <sub>11</sub> ] <sup>2-</sup>                                       | -569.6334Eh              | -569.6324Eh  | 0.0414Eh    | -569.6738Eh           |
| [GeB <sub>11</sub> H <sub>11</sub> ] <sup>2-</sup>                                       | -2357.3931Eh             | -2357.3922Eh | 0.0428Eh    | -2357.4350Eh          |
| [SnB <sub>11</sub> H <sub>11</sub> ] <sup>2-</sup>                                       | -494.5461Eh              | -494.5452Eh  | 0.0440Eh    | -494.5891Eh           |
| [PbB <sub>11</sub> H <sub>11</sub> ] <sup>2-</sup>                                       | -473.1383Eh              | -473.1374Eh  | 0.0451Eh    | -473.1824Eh           |
| [(CB <sub>11</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>2-</sup>                        | -636.4633Eh              | -636.4623Eh  | 0.0562Eh    | -636.5185Eh           |
| [(SiB <sub>11</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>2-</sup>                       | -1139.3379Eh             | -1139.3369Eh | 0.0621Eh    | -1139.3989Eh          |
| [(GeB <sub>11</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>2-</sup>                       | -4714.8216Eh             | -4714.8207Eh | 0.0649Eh    | -4714.8855Eh          |
| [(SnB <sub>11</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>2-</sup>                       | -989.0885Eh              | -989.0875Eh  | 0.0676Eh    | -989.1551Eh           |
| [(SnB <sub>11</sub> H <sub>10</sub> )(SnB <sub>11</sub> H <sub>11</sub> )] <sup>3-</sup> | -988.4279Eh              | -988.4276Eh  | 0.0648Eh    | -988.4924Eh           |
| [(PbB <sub>11</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>2-</sup>                       | -946.2364Eh              | -946.2354Eh  | 0.0708Eh    | -946.3063Eh           |

The energy of the reactions for the dimers are given in Table 4. This energy is derived from the subtracting the total thermal energy (U) of the product and 2 mol of reactants, then the Hartree value is converted to kcal/mol.

Table 4. DFT calculated Energy of the reaction using total thermal energy (U), enthalpy (H), Gibbs free energy (G) and entropy (S) for group 14 hetero-*closo*-dodecaborate dimers at BP86/Def2-TZVP level.

| Dimers | $\Delta U$ |          | $\Delta H$ |          | $\Delta G$ |          | $\Delta S$ |          |
|--------|------------|----------|------------|----------|------------|----------|------------|----------|
|        | Hartrees   | kcal/mol | Hartrees   | kcal/mol | Hartrees   | kcal/mol | Hartrees   | Kcal/mol |
| 1      | 1.2765     | 801.0    | 1.2755     | 800.4    | 1.2989     | 815.1    | -0.0232    | -14.6    |
| 2      | -0.0710    | -44.6    | -0.0721    | -45.2    | -0.0513    | -32.2    | -0.0207    | -13.0    |
| 3      | -0.0354    | -22.2    | -0.0363    | -22.8    | -0.0155    | -9.7     | -0.0207    | -13.0    |
| 4A     | 0.0037     | 2.4      | 0.0029     | 1.8      | 0.0231     | 14.5     | -0.0204    | -12.8    |
| 4B     | 0.6643     | 416.9    | 0.6628     | 415.9    | 0.6858     | 430.3    | -0.0232    | -14.6    |
| 5      | -0.0401    | 25.2     | 0.0394     | 24.7     | 0.0585     | 36.7     | -0.0194    | -12.2    |

Energy of the reaction by using total thermal energy (U),

$$\Delta U = U_{\text{Product}} - U_{\text{Reactants}} \quad 9$$

Enthalpy of the reaction,

$$\Delta H = H_{\text{Product}} - H_{\text{Reactants}} \quad 10$$

Gibbs free energy,

$$\Delta G = \Delta H - (T \times \Delta S) \quad 11$$

Gibbs free energy of the reaction,

$$\Delta G = G_{\text{Product}} - G_{\text{Reactants}}$$

The borates of silicon and germanium have the energies using total thermal energy is  $-44.5531$  kcal/mol and  $-22.2138$  kcal/mol respectively, which indicates that the reactions are feasible in which germa-*closo*-dodecaborate dimer is an experimentally known cluster. The energy of the reaction for the formation of diplumba-*closo*-dodecaborate is  $25.2$  kcal/mol. Interestingly for the stanna-*closo*-dodecaborate dimer with Sn=Sn (**4A**) and Sn-B (**4B**) has  $2.3835$  kcal/mol and  $416.9$  kcal/mol. The cluster (**4A**) has  $2.3$  kcal/mol which is lower energy than an experimentally known cluster hetero-*closo*-distannaborate (**4B**) with Sn-B bond [44] and confirming the viability of the modelled cluster hetero-*closo*-distannaborate (**4A**) with Sn=Sn bonding.

The change in entropy of the reaction, has been calculated using equation (12)



Two moles of reactant  $\rightarrow$  1 mol of dimer

$$\Delta S = S_{\text{Product}} - S_{\text{Reactant}}$$

The DFT (BP86/Def2-TZVP) computed entropy of the reaction values for the experimentally known dimers (**3**) and (**4B**) are  $-0.0207$  Eh and  $-0.0232$  Eh respectively, Table 4. The plumba-*closo*-dodecaborate dimer (**5**) has the slightly higher value  $-0.0194$  Hartrees than the remaining dimers. The change in Gibbs free energy for the dimer forming reaction confirm the more feasible nature of dimers **2**, **3**, **4A** and **5** and less feasible nature of **1** with  $1.2989$  Hartrees. From the free energy values of the dimer forming reaction, the following order has been depicted for the formation of dimer,  $2 > 3 > 4A > 5 > 4B$ .

### 3.8. Spectroscopic properties

The hetero-*closo*-dodecaborate has three types of protons and borons which lead to three types of chemical shifts ( $\delta$ ), upper belt (5B (1–5) and 5H (1–5)), lower belt (5B (6–10) and H (6–10)) and apical vertex (B11 and H11). The DFT computed chemical shift values are given in Tables S10–S14. The  $^{11}\text{B}$  nmr chemical shifts are resonates in 1:5:5 ratio. Usually upper belt borons and protons has lower chemical shift than lower belt and apical position through the higher shielding effects presents in the upper belt than in the remaining atoms. According to the presence of group-14 heteroatoms, the upper belt obtains higher shielding effect. The difference between the lower belt and upper belt is around 2–4 ppm. When comparing these two belts with apical vertex, the chemical shift of apical position is very high. The apical vertex B and H atoms are highly deshielded, these elements utilize higher resonating energy around 4–10 ppm than the remaining atoms. The DFT (BP86/Def2-TZVP) computed chemical shift for C atom of cluster (**1a**) is  $58.74$  which is close to the experimental value of  $51.5$  ppm. An upper belt boron resonates around  $-21.22$  to  $-21.26$  ppm which is close to the experimental value of  $-16.31$  ppm. The lower belt boron resonates around  $-17.24$  to  $-17.27$  ppm which is also close to the experimental value of  $-13.30$  ppm. The apical boron B11 [DFT  $-11.21$  ppm; Experimental  $-6.90$  ppm]. Is less shielded when compared to those of upper best and lower belt boron atoms. The  $^1\text{H}$  nmr shows that the chemical shift values of hydrogen atoms are in 1:5:5 ratio, the proton connected with the C atom resonates at  $2.17$  ppm which is close to the experimental value of  $2.24$  ppm [85,86]. Similarly, each monomers (**2a–5a**) and dimers (**1–5**) except (**4B**) computed chemical shifts of  $^{11}\text{B}$ ,  $^1\text{H}$  resonates like 1:5:5 for monomers and both cages of dimers. The  $^{13}\text{C}$  nmr of carborate dimer  $[(\text{CB}_{11}\text{H}_{11})_2]^{2-}$  (**1**) resonates around  $86.0$  ppm. The silicon atom of the methyl substituted sila-*closo*-dodecaborane (**2b**) The  $^{29}\text{Si}$  nmr of experimentally known cluster (**2b**) has been computed at DFT (BP86/Def2-TZVP) level as  $-36.84$  ppm which is close to the experimental value of  $-28.0$  ppm [2,20,73]. The  $[(\text{GeB}_{11}\text{H}_{11})_2]^{2-}$  (**3**) cluster resonates 1:5:5 ratio [44]. In the stannaborate (**4a**) and plumbaborate (**5a**) monomers the  $^{11}\text{B}$  nmr shows peaks in the 1:5:5 ratio. The DFT and experimental values are  $-8.72$  [5.1]:  $16.23$  [ $-10.6$ ]:  $18.14$  [ $-11.9$ ] ppm for (**4a**) and  $-7.74$  [ $-2.9$ ]:  $18.53$  [ $-10.3$ ]:  $14.46$  [ $-5.0$ ] ppm for (**5a**) [20], whereas the dimers resonates at  $-14.0$ :  $21.0$ :  $19.0$  and  $-21.0$ :  $24.0$ :  $17.0$  (B11:B6-B10:B1-B5). The boron atoms of  $(\text{SnB}_{11}\text{H}_{10})(\text{SnB}_{11}\text{H}_{11})^{3-}$  (**4B**) cluster resonates at  $-11.36$  [ $-13.6$ ] (B1);  $-17.59$  [ $-8.0$ ] (B8);  $-8.57$  [ $-3.6$ ] (B11);  $-14.83$  [ $-10.0$ ] (B2),  $-14.92$  [ $-10$ ] (B5);  $-14.92$  [ $-8.6$ ] (B3),  $-14.92$  [ $-8.6$ ] (B4);  $-17.60$  [ $-11.4$ ] (B6),  $-17.44$  [ $-11.4$ ] (B10),  $-17.58$  [ $-11.4$ ] (B7),  $-17.41$  [ $-11.4$ ] (B9);  $-21.91$  to  $-22.50$  [ $-15.8$ ] (B12-B22) ppm. An experimentally reported  $^{11}\text{B}$  nmr ratio is 1:1:1:2:2:2:11. But, the ratio of cage **2** in (**4B**) normal icosahedrons 1:5:5 ratio peakas

-21.19 (B1-B5): 22.51 (B6-B10): 15.36 (B11) ppm. The apical vertex H11 resonates at 2.447 ppm which is less shielded as expected, than the  $^1\text{H}$  nmr chemical shifts of the remaining hydrogen atoms of  $(\text{SnB}_{11}\text{H}_{10})(\text{SnB}_{11}\text{H}_{11})]^{3-}$  (**4B**) cluster [44]. The DFT (BP86/Def2-TZVP) calculated  $^{11}\text{B}$ ,  $^{13}\text{C}$ ,  $^1\text{H}$  and  $^{29}\text{Si}$  nmr chemical shift values are helpful in assigning the individual atoms of the monomeric and dimeric clusters.

## 4. Conclusions

The bonding interactions of hetero-*closo*-dodecaborates (**1a** – **5a**) and their dimers (**1**–**5**) are analyzed using density functional theory (DFT) and DFT-D3 method at BP86/Def2-TZVP level of theory. The viable nature of the dimeric hetero-*closo*-dodecaborate clusters (**2**–**5**) are confirmed through DFT computations. Although, normal DFT optimization leads to the closer values towards experimentally obtained bond parameters, inclusion of dispersion correlation suggested by Grimme et al. improves the metrical parameters significantly ( $\sim 0.04\text{\AA}$  in bond length) towards an experimental value. Further, inclusion of dispersion correction provides the energies more accurately. The DFT computed global and local descriptors have predict correctly the labile nature of the stannaborate dimer (**4A**) with 'Sn=Sn' double bond. Both DFT and experimental studies confirm the more stable nature of the stannaborate dimer (**4B**) with possible Sn-B bonding interaction. The global electrophilicity index ( $\omega$ ) of the monomeric clusters (**2a**, **2**, **3a**, **4a** and **5a**) are higher than the dimers except carborates. DFT (BP86/Def2-TZVP) calculations on the clusters confirm the thermal stability of the newly modelled dimeric clusters (**2**–**5**) which could have diverse applications. ELF analysis confirm the high electron density around, 0.5 to 0.9 range for E=E bond in each dimers except the dimer (**5**). From the free energy values of the dimer forming reaction, the following order, **2** > **3** > **4A** > **5** > **4B**., has been depicted for the formation of dimer. The DFT (BP86/Def2-TZVP) computed spectroscopic values are in good agreement with the experimental values when available or with similar clusters. The  $^{13}\text{C}$ ,  $^1\text{H}$ ,  $^{29}\text{Si}$ , and  $^{11}\text{B}$  nmr chemical shift values computed at DFT (BP86/Def2-TZVP) level are highly useful in identifying and assigning the individual atoms of the monomeric and dimeric clusters.

## CRedit authorship contribution statement

**Brindha Veerappan:** Writing – original draft, Methodology, Investigation, Data curation. **Krishnamoorthy Bellie Sundaram:** Writing – review & editing, Supervision, Investigation, Conceptualization.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Appendix A. Supplementary data

The following is the Supplementary data to this article:

 [Download: Download Word document \(4MB\)](#)

Multimedia component 1.

[Recommended articles](#)

## Data availability

No data was used for the research described in the article.

## References

- [1] P.P. Power  
 $\pi$ -Bonding and the lone pair effect in multiple Bonds between Heavier Main Group elements  
Chem. Rev., 99 (1999), pp. 3463-3504, [10.1021/cr9408989](https://doi.org/10.1021/cr9408989) ↗

[Google Scholar](#) ↗

- [2] K.W. Klinkhammer, W. Schwarz  
Bis(hypersilyl)tin and bis(hypersilyl)lead, two electron-rich carbene homologs  
Angew. Chem., Int. Ed. Engl., 134 (1995), pp. 1334-1336, [10.1002/anie.199513341](#) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [3] H. Zhiyuan, L. Lingyu, J.D.Z. Felix, X. Xiaolian, W.E. Andreas, Y. KaKing, D. Serhiy, V. Jarl Ivar van der, B. Bas, K. Jeremy  
Reactivity of a unique Si(I)–Si(I)–Based $\eta^2$ -Bis(silylene) iron complex  
Inorg. Chem., 61 (30) (2022), pp. 11725-11733, [10.1021/acs.inorgchem.2c01369](#) ↗  
[Google Scholar](#) ↗
- [4] M. Stürmann, W. Saak, K.W. Klinkhammer, M. Weidenbruch  
A Heteroleptic (E)-1,2-Diaryl-1,2-disilyl distannene without Donor Stabilization, Zeitschrift. Für.  
Anorganische. Und  
Allgemeine. Chemie., 625 (1999), pp. 1955-1956, [10.1002/\(SICI\)1521-3749\(199912\)625:12<1955::AID-ZAAC1955>3.0.CO;2-J](#) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [5] R. Bashkurov, N. Fridman, D. Bravo-Zhivotovskii, Y. Apeloig  
The first planar, not twisted, Distannene –A structural alkene analog, synthesis, isolation and x-  
ray crystallography characterization  
Chem. Eur J., 29 (69) (2023), Article e202302678, [10.1002/chem.202302678](#) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [6] P.J. Davidson, D.H. Harris, M.F. Lappert  
Subvalent Group 4B metal alkyls and amides. Part I. The synthesis and physical properties of  
kinetically stable Bis[bis(trimethylsilyl)methyl]-germanium(II), -tin(II), and -lead(II)  
J. Chem. Soc. Dalton Trans. (1976), pp. 2268-2274, [10.1039/DT9760002268](#) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [7] K.W. Klinkhammer, T.F. Fässler, H. Grützmacher  
The Formation of HeterolepticCarbene homologues by Ligand exchange - synthesis of the first  
plumbanediyl dimer  
Angew. Chem. Int. Ed., 37 (1998), pp. 124-126, [10.1002/\(SICI\)1521-3773\(19980202\)37:1/2<124::AID-ANIE124>3.0.CO;2-C](#) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [8] S. Wang, H.J. Li, T.S. Kuo, L.C. Shen, H.J. Liu  
Ambiphilic Nature of dipyrrolylpyridine - supported divalent germanium and tin compounds  
Organometallics, 40 (2021), pp. 3659-3667, [10.1021/acs.organomet.1c00494](#) ↗  
[View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [9] S. Nagase  
Multiple bonds between lead atoms and short bonds between transition metals  
Pure Appl. Chem., 85 (2012), pp. 649-659, [10.1351/PAC-CON-12-08-04](#) ↗  
[Google Scholar](#) ↗
- [10] J.D. Guo, D.J. Liptrot, S. Nagase, P.P. Power  
The multiple bonding in heavier group 14 element alkene analogues is stabilized mainly by  
dispersion force effects  
Chem. Sci., 6 (2015), pp. 6235-6244, [10.1039/C5SC02707A](#) ↗  
[View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [11] S. Körbe, P.J. Schreiber, J. Michl  
Chemistry of the carba-*closo*-dodecaborate(–) anion, CB11<sup>H</sup>12<sup>–</sup>  
Chem. Rev., 106 (2006), pp. 5208-5249, [10.1021/cr050548u](#) ↗

- [View in Scopus ↗](#) [Google Scholar ↗](#)
- [12] T. Gädt, L. Wesemann  
Stanna-*closo*-dodecaborate chemistry  
Organometallics, 26 (2007), pp. 2474-2481, [10.1021/om061042k ↗](#)  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [13] A. Stock, O. Maesenez  
Reports of the German Chemical Society, 45 (3) (1912), pp. 3539-3568, [10.1002/cber.191204503113 ↗](#)  
boron hydrogens  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [14] W.N. Lipscomb  
Boron Hydrides, the Physical Inorganic Chemistry Series  
Harvard University, New York (1963)  
[Google Scholar ↗](#)
- [15] W.N. Lipscomb  
Bonding in boron hydrides  
Pure Appl. Chem., 29 (4) (1972), pp. 493-512, [10.1351/pac197229040493 ↗](#)  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [16] N.S. Hosmane, J.A. Maguire  
Boron Hydrides, based in part on the article Boron Hydrides by James T. Spencer which appeared in the *Encyclopedia of Inorganic Chemistry*  
Encyclopedia of Inorganic Chemistry (first ed.), John Wiley & Sons, Ltd. (2006), [10.1002/0470862106.ia024 ↗](#)  
ISBN: 9780470862100  
[Google Scholar ↗](#)
- [17] I.E. Golub, O.A. Filippov, V.A. Kulikova, N.V. Belkova, L.M. Epstein, E.S. Shubina  
Thermodynamic hydricity of small borane clusters and polyhedral *closo*-Boranes  
Molecules, 25 (12) (2020), p. 2920, [10.3390/molecules25122920 ↗](#)  
[Google Scholar ↗](#)
- [18] J. Aihara  
Three-dimensional aromaticity of polyhedral boranes  
J. Am. Chem. Soc., 100 (11) (1978), pp. 3339-3342  
[https://pubs.acs.org/doi/10.1021/ja00479a015 ↗](https://pubs.acs.org/doi/10.1021/ja00479a015)  
[Crossref ↗](#) [View in Scopus ↗](#) [Google Scholar ↗](#)
- [19] R.B. King  
Three-Dimensional aromaticity in polyhedral boranes and related molecules  
Chem. Rev., 101 (5) (2001), pp. 1119-1152, [10.1021/cr000442t ↗](#)  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [20] R.W. Chapman, J.G. Kester, K. Folting, W.E. Streib, L.J. Todd  
Synthesis and chemistry of B<sub>11</sub>H<sub>11</sub>Sn<sup>2-</sup> and its germanium and lead analogs. Crystal structure of [B<sub>11</sub>H<sub>11</sub>SnCH<sub>3</sub>]PPh<sub>3</sub>CH<sub>3</sub>.CH<sub>2</sub>Cl<sub>2</sub>  
Inorg. Chem., 31 (6) (1992), pp. 979-983, [10.1021/ic00032a011 ↗](#)  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [21] T. Wutz, F. Diab, L. Wesemann  
Synthesis and coordination chemistry of a P-Sn-chelating ligand based on stanna-*closo*-dodecaborate  
Eur. J. Inorg. Chem., 2017 (2017), pp. 4645-4652, [10.1002/ejic.201700471 ↗](#)

[View in Scopus](#) [Google Scholar](#)

- [22] M.A. Fox, T.B. Marder, L. Wesemann  
DFT studies of the  $\sigma$ -donor/ $\pi$ -acceptor properties of  $[\text{SnCB}_{10}\text{H}_{11}]^-$  and its relationship to  $[\text{SnCl}_3]^-$ , CO, PF<sub>3</sub>,  $[\text{SnB}_{11}\text{H}_{11}]^{2-}$ , SnC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>, and related SnC<sub>2</sub>B<sub>n</sub>H<sub>n+2</sub> compounds  
Can. J. Chem., 87 (1) (2009), pp. 63-71, [10.1139/v08-081](#)  
[View in Scopus](#) [Google Scholar](#)
- [23] M.F. Hawthorne, A. Maderna  
Applications of radiolabeled boron clusters to the diagnosis and treatment of cancer  
Chem. Rev., 99 (1999), pp. 3421-3434, [10.1021/cr000001+](#)  
Published on the Web Nov. 13, 1999. Chem Rev 100(3) (2000) 1165  
[View in Scopus](#) [Google Scholar](#)
- [24] M.F. Hawthorne, S.S. Jalisatgi, A.V. Safronov, H.B. Lee, J. Wu  
Chemical Hydrogen Storage Using Polyhedral Borane Anions and aluminum-ammonia-borane Complexes  
International Institute of Nano and Molecular Medicine, University of, Missouri, United States (2010), [10.2172/990217](#)  
[Google Scholar](#)
- [25] W.H. Jin, C. Seldon, M. Butkus, W. Sauerwein, H.B. Giap  
A review of Boron Neutron capture therapy: its history and Current challenges  
International Journal of Particle Therapy, 9 (2022), pp. 71-82, [10.14338/IJPT-22-00002.1](#)  
 [View PDF](#) [View article](#) [View in Scopus](#) [Google Scholar](#)
- [26] L. Pazderová, E.Z. Tüzün, D. Baval, M. Litecká, L. Fojt, B. Grúner  
Chemistry of carbon-substituted derivatives of cobalt Bis(dicarbollide)(1-) ion and recent progress in boron substitution  
Molecules, 28 (2023), p. 6971, [10.3390/molecules28196971](#)  
[View in Scopus](#) [Google Scholar](#)
- [27] S.H. Payandeh, R. Asakura, P. Avramidou, D. Rentsch, Z. Łodziana, R. Černý, A. Remhof, C. Battaglia  
Nido-Borate/Closo-Borate mixed-anion electrolytes for All-SolidState batteries  
Chem. Mater., 32 (2020), pp. 1101-1110, [10.1021/acs.chemmater.9b03933](#)  
[View in Scopus](#) [Google Scholar](#)
- [28] T.A. Hales, K.T. Moller, T.D. Humphries, A.M. D'Angelo, C.E. Buckley, M. Paskevicius  
Investigating the potential of alkali metal plumba-closo-dodecaborate (B<sub>11</sub>H<sub>11</sub>Pb)<sup>2-</sup> salts as solid-state battery electrolytes  
J. Phys. Chem. C, 127 (2023), pp. 949-957, [10.1021/acs.jpcc.2c07226?urlappend=%3Fref%3DPDF&jav=VoR&rel=cite-as](#)  
[View in Scopus](#) [Google Scholar](#)
- [29] Z. Lu, F. Ciucci  
Metal borohydrides as electrolytes for solid-state Li, Na, Mg, and Ca batteries: a first-principles Study  
Chem. Mater., 29 (2017), pp. 9308-9319  
<https://pubs.acs.org/doi/10.1021/acs.chemmater.7b03284>  
[Crossref](#) [View in Scopus](#) [Google Scholar](#)
- [30] A. Avelar, F.S. Tham, C.A. Reed  
Superacidity of boron acids H<sub>2</sub>(B<sub>12</sub>X<sub>12</sub>) (X=Cl, Br)  
Angew. Chem., Int. Ed. Engl., 48 (19) (2009), pp. 3491-3493, [10.1002/anie.200900214](#)  
[View in Scopus](#) [Google Scholar](#)
- [31] B. Gruner, J. Rais, P. Selucky, M. Lucanikova

## Recent progress in extraction agents based on cobalt Bis(Dicarbollides) for partitioning of radionuclides from high-level nuclear waste

Boron Science: New Technologies and Applications, N. S. Hosmane (2011), pp. 463-490, [10.1201/b11199](https://doi.org/10.1201/b11199) ↗  
eISBN9780429062735

[Google Scholar](#) ↗

- [32] J. Fanfrlik, J. Brynda, J. Rezac, P. Hobza, M. Lepsik  
Interpretation of Protein/Ligand crystal structure using QM/MM calculations: case of HIV-1 Protease/Metallacarborane complex  
J. Phys. Chem. B, 112 (2008), pp. 15094-15102, [10.1021/jp803528w](https://doi.org/10.1021/jp803528w) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [33] R. Nnez, M. Tarres, A. Ferrer-Ugalde, F. FabriziDebiani, F. Teixidor  
Electrochemistry and photoluminescence of icosahedral carboranes, boranes, metallacarboranes, and their derivatives  
Chem. Rev., 116 (23) (2016), pp. 14307-14378, [10.1021/acs.chemrev.6b00198](https://doi.org/10.1021/acs.chemrev.6b00198) ↗  
[Google Scholar](#) ↗
- [34] R.K. Roy, S. Krishnamurti, P. Geerlings, S. Pal  
Local softness and hardness based reactivity descriptors for predicting Intra – and intermolecular reactivity sequences: carbonyl compounds  
J. Phys. Chem., 102 (1998), pp. 3746-3755, [10.1021/jp973450v](https://doi.org/10.1021/jp973450v) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [35] A. Aizman, R. Contreras, P. Pérez  
Relationship between local electrophilicity and rate coefficients for the hydrolysis of carbenium ions  
Tetrahedron, 61 (2005), pp. 889-895, [10.1016/j.tet.2004.11.014](https://doi.org/10.1016/j.tet.2004.11.014) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [36] C. Morell, A. Grand, A. Toro-Labbé  
Theoretical support for using the  $\Delta f(r)$  descriptor  
Chem. Phys. Lett., 425 (2006), pp. 342-346, [10.1016/j.cplett.2006.05.003](https://doi.org/10.1016/j.cplett.2006.05.003) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [37] C. Cárdenas, N. Rabi, P.W. Ayers, C. Morell, P. Jaramillo, P. Fuentealba  
Chemical reactivity descriptors for ambiphilic reagents: dual descriptor, local hypersoftness and electrostatic potential  
J. Phys. Chem., 113 (2009), pp. 8660-8667, [10.1021/jp902792n](https://doi.org/10.1021/jp902792n) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [38] J. Padmanabhan, R. Parthasarathi, M. Elango, V. Subramanian, B.S. Krishnamoorthy, S. Gutierrez-Oliva, A. Toro-Labbe, D.R. Roy, P.K. Chattaraj  
Multiphilic descriptor for chemical reactivity and selectivity  
J. Phys. Chem., 111 (2007), pp. 9130-9138, [10.1021/jp0718909](https://doi.org/10.1021/jp0718909) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [39] C. Morell, A. Grand, A. Toro-Labbé  
New dual descriptor for chemical reactivity  
J. Phys. Chem., 109 (2005), pp. 205-212, [10.1021/jp046577a](https://doi.org/10.1021/jp046577a) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [40] G.I. Cárdenas-Jirón, S. Gutiérrez-Oliva, J. Melin, A. Toro-Labbé  
Relations between potential energy, electronic chemical potential and hardness profiles

- J. Phys. Chem., 101 (1997), pp. 4621-4627, [10.1021/jp9638705](https://doi.org/10.1021/jp9638705) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [41] R.S. Mulliken  
Molecular compounds and their spectra II  
J. American Chem. Soc., 74 (3) (1952), pp. 811-824, [10.1021/ja01123a067](https://doi.org/10.1021/ja01123a067) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [42] T. Kar, S. Scheiner, A.B. Sannigrahi  
Ab initio calculations of hardness and chemical potential of open shell systems using SCF, MP2 and MP4 methods  
J. Mol. Struct.: THEOCHEM, 427 (1998), pp. 79-85, [10.1016/S0166-1280\(97\)00172-3](https://doi.org/10.1016/S0166-1280(97)00172-3) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [43] J. Luo, Z.Q. Xue, W.M. Liu, J.L. Wu, Z.Q. Yang  
Koopmans' Theorem for large molecular systems within density functional theory  
J. Phys. Chem., 110 (2006), pp. 12005-12009, [10.1021/jp063669m](https://doi.org/10.1021/jp063669m) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [44] T. Gädt, J.-A. Dimmer, S. Fleischhauer, A. Frank, C. Nickl, T. Wutz, K. Eichele, L. Wesemann  
Oxidation of germa- and stanna-*closo*-dodecaborate  
Dalton Trans., 44 (2015), pp. 4726-4731, [10.1039/C5DT00099H](https://doi.org/10.1039/C5DT00099H) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [45] V. Brindha, B.S. Krishnamoorthy  
Isomer preferences and structural studies on cobaltaboranes – a theoretical investigation  
BiointerfaceReseach in Applied Chemistry, 14 (3) (2024), p. 73, [10.33263/BRIAC143.073](https://doi.org/10.33263/BRIAC143.073) ↗  
[Google Scholar](#) ↗
- [46] V. Nagalakshmi, R. Nandhini, V. Brindha, B.S. Krishnamoorthy, K. Balasubramani  
Half-sandwich ruthenium(II) complexes containing biphenylamine based Schiff base ligands: synthesis, structure and catalytic activity in amidation of various aldehydes  
J. Organomet. Chem., 912 (2020), Article 121175, [10.1016/j.jorgchem.2020.121175](https://doi.org/10.1016/j.jorgchem.2020.121175) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [47] S. Gomathi, K. Kavitha, N. Savitha, A. Mohana, V. Brindha, B.S. Krishnamoorthy  
Molecular structure, reactivity and spectroscopic properties of Hallucinogens Psilocybin, mescaline and their Derivatives–A computational Study  
Lett. Appl. Nano. Bio. Sci., 12 (2022), p. 105, [10.33263/LIANBS124.105](https://doi.org/10.33263/LIANBS124.105) ↗  
[Google Scholar](#) ↗
- [48] Yusuf Sert, Mustafa R. Albayati, Fatih Şen, Necmi Dege  
The DFT and in-silico analysis of 2,2'-((1e,1'e)-((3,3'-dimethyl-[1,1'-biphenyl]-4,4'-diyl)bis(azanylylidene))bis(methanylylidene))diphenol molecule  
Colloids Surf. A Physicochem. Eng. Asp., 687 (2024), Article 133444, [10.1016/j.colsurfa.2024.133444](https://doi.org/10.1016/j.colsurfa.2024.133444) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [49] Necmi Dege, Halil Gökce, Onur Erman Doğan, Gökhan Alpaslan, Tuğgan Açar, S. Muthu, Yusuf Sert  
Quantum computational, spectroscopic investigations on N-(2-((2-chloro-4,5-dicyanophenyl)amino)ethyl)-4-methylbenzenesulfonamide by DFT/TD-DFT with different solvents, molecular docking and drug-likeness researches  
Colloids Surf. A Physicochem. Eng. Asp., 638 (2022), Article 128311, [10.1016/j.colsurfa.2022.128311](https://doi.org/10.1016/j.colsurfa.2022.128311) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [50] Mahmood A. Albo Hay Allah, Asim A. Balakit, Hamida Idan Salman, Ahmed Abdulridha Ali, Yusuf Sert

New heterocyclic compound as carbon steel corrosion inhibitor in 1 M H<sub>2</sub>SO<sub>4</sub>, high efficiency at low concentration: experimental and theoretical studies

J. Adhes. Sci. Technol., 37 (3) (2022), pp. 525-547, [10.1080/01694243.2022.2034588](https://doi.org/10.1080/01694243.2022.2034588) ↗

[Google Scholar](#) ↗

[51]

F. Neese

Software update: the ORCA program system—Version 5.0

WIREs Comput. Mol. Sci., 12 (2022), Article e1606, [10.1002/wcms.1606](https://doi.org/10.1002/wcms.1606) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[52]

S.H. Vosko, L. Wilk, M. Nusair

Accurate spin-dependent electron liquid correlation energies for local spin density calculations: a critical analysis

Can. J. Phys., 58 (1980), pp. 1200-1211, [10.1139/p80-159](https://doi.org/10.1139/p80-159) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[53]

A.D. Becke

Density functional calculations of molecular bond energies

J. Chem. Phys., 84 (1986), pp. 4524-4529, [10.1063/1.450025](https://doi.org/10.1063/1.450025) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[54]

A.D. Becke

Density- functional exchange-energy approximation with correct asymptotic behavior

Phys. Rev., 38 (1988), pp. 3098-3100, [10.1103/PhysRevA.38.3098](https://doi.org/10.1103/PhysRevA.38.3098) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[55]

J.P. Perdew

Density-functional approximation for the correlation energy of the inhomogeneous electron gas

Phys. Rev. B, 33 (1986), pp. 8822-8824, [10.1103/PhysRevB.33.8822](https://doi.org/10.1103/PhysRevB.33.8822) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[56]

B.S. Krishnamoorthy, A. Thakur, K.K.V. Chakrahari, S.K. Bose, P. Hamon, T. Roisnel, S. Kahlal, S. Ghosh, J.F. Halet

Theoretical and experimental investigations on hypoelectronic heterodimetallaboranes of Group 6 transition metals

Inorg. Chem., 51 (2012), pp. 10375-10383, [10.1021/ic301571e](https://doi.org/10.1021/ic301571e) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[57]

B.S. Krishnamoorthy, S. Kahlal, S. Ghosh, J.F. Halet

Electronic, geometrical and thermochemical studies on group-14 element-diruthenaborane cluster compounds: a theoretical investigation

Theor. Chem. Acc., 132 (2013), p. 1356, [10.1007/s00214-013-1356-6](https://doi.org/10.1007/s00214-013-1356-6) ↗

[Google Scholar](#) ↗

[58]

K. Geetharani, B.S. Krishnamoorthy, S. Kahlal, S.M. Mobin, S. Ghosh, J.F. Halet

Synthesis and characterization of HypoelectronicTantalaboranes: comparison of geometric and electronic structures of [(Cp\*TaX)<sub>2</sub>B<sub>5</sub>H<sub>11</sub>] (X = Cl, Br, I)

Inorg. Chem., 51 (2012), pp. 10176-10184, [10.1021/ic300848f](https://doi.org/10.1021/ic300848f) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

[59]

F. Weigend, R. Ahlrichs

Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: design and assessment of accuracy

Phys. Chem. Chem. Phys., 7 (2005), p. 3297, [10.1039/b508541a](https://doi.org/10.1039/b508541a) ↗

[View in Scopus](#) ↗ [Google Scholar](#) ↗

- [60] S. Grimme, J. Antony, S. Ehrlich, H. Krieg  
A consistent and accurate *ab initio* parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu  
J. Chem. Phys., 132 (2010), Article 154104, [10.1063/1.3382344](https://doi.org/10.1063/1.3382344) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [61] S. Grimme, S. Ehrlich, L. Goerigk  
Effect of the damping function in dispersion corrected density functional theory  
J. Comput. Chem., 32 (2011), pp. 1456-1465, [10.1002/jcc.21759](https://doi.org/10.1002/jcc.21759) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [62] E.R. Johnson, A.D. Becke  
A post-hartree-fock model of intermolecular interactions: inclusion of higher-order corrections  
J. Chem. Phys., 124 (2006), Article 174104, [10.1063/1.2190220](https://doi.org/10.1063/1.2190220) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [63] E.R. Johnson, A.D. Becke  
A post-hartree-fock model of intermolecular interactions  
J. Chem. Phys., 123 (2005), Article 024101, [10.1063/1.1949201](https://doi.org/10.1063/1.1949201) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [64] J. Mareš, J. Vaara  
*Ab initio* paramagnetic NMR shifts *via* point-dipole approximation in a large magnetic-anisotropy Co(II) complex  
Phys. Chem. Chem. Phys., 20 (2018), pp. 22547-22555, [10.1039/C8CP04123G](https://doi.org/10.1039/C8CP04123G) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [65] Z.S. Safi, N. Wazzan  
DFT calculations of  $^1\text{H}$  and  $^{13}\text{C}$ -NMR chemical shifts of 3-methyl-1-phenyl-4-(phenyldiazenyl)-1H-pyrazol-5-amine in solution  
Sci. Rep., 12 (2022), Article 17798, [10.1038/s41598-022-22900-y](https://doi.org/10.1038/s41598-022-22900-y) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [66] E.D. Becker  
High Resolution NMR: Theory and Chemical Applications  
(third ed.) (2000), [10.1016/B978-0-12-084662-7.X5044-3](https://doi.org/10.1016/B978-0-12-084662-7.X5044-3) ↗  
ISBN: 9780120846627  
[Google Scholar ↗](#)
- [67] P.K. Chattaraj, B. Maiti, U. Sarkar  
Philicity: a unified treatment of chemical reactivity and selectivity  
J. Phys. Chem., 107 (2003), pp. 4973-4975, [10.1021/jp034707u](https://doi.org/10.1021/jp034707u) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [68] D.R. Roy, R. Parthasarathi, J. Padmanabhan, U. Sarkar, V. Subramanian, P.K. Chattaraj  
Careful scrutiny of the philicity concept  
J. Phys. Chem., 110 (2006), pp. 1084-1093, [10.1021/jp053641v](https://doi.org/10.1021/jp053641v) ↗  
[View in Scopus ↗](#) [Google Scholar ↗](#)
- [69] W. Tiznado, O.B. Oña, V.E. Bazterra, M.C. Caputo, J.C. Facelli, M.B. Ferraro, P. Fuentealba  
Theoretical study of the adsorption of H on  $\text{Si}_n$  clusters ( $n= 3-10$ )  
J. Chem. Phys., 123 (2005), Article 214301, [10.1063/1.2128675](https://doi.org/10.1063/1.2128675) ↗  
[Google Scholar ↗](#)

- [70] E. Osorio, M.B. Ferraro, O.B. Oña, C. Cardenas, P. Fuentealba, W. Tiznado  
Assembling small silicon clusters using criteria of maximum matching of the Fukui functions  
J. Chem. Theor. Comput., 7 (12) (2011), pp. 3995-4001, [10.1021/ct200643z](https://doi.org/10.1021/ct200643z) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [71] O. Yañez, R. Báez-Grez, D. Inostroza, R. Pino-Rios, W.A. Rabanal-León, J. Contreras-García, C. Cardenas, W. Tiznado  
Kick–Fukui: a Fukui function-guided method for molecular structure prediction  
J. Chem. Inf. Model., 61 (8) (2021), pp. 3955-3963, [10.1021/acs.jcim.1c00605](https://doi.org/10.1021/acs.jcim.1c00605) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [72] E. Osorio, A.P. Sergeeva, J.C. Santos, W. Tiznado  
Theoretical study of the  $\text{Si}_{5-n}(\text{BH})_n^{2-}$  and  $\text{Na}(\text{Si}_{5-n}(\text{BH})_n)^-$  ( $n = 0-5$ ) systems  
Phys. Chem. Chem. Phys., 14 (2012), pp. 16326-16330, [10.1039/C2CP42674A](https://doi.org/10.1039/C2CP42674A) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [73] Lars Wesemann  
UlliEnglert, the first *closo*-monosilaborane  
Angew. Chem., 108 (1996), pp. 586-587, [10.1002/anie.199605271](https://doi.org/10.1002/anie.199605271) ↗  
[Google Scholar](#) ↗
- [74] Torben Gadt, Lars Wesemann  
Stanna-*closo*-dodecaborate: the Crystal Structure of  $[\text{Li}(\text{thf})_3]_2[\text{SnB}_{11}\text{H}_{11}]$ , Vibrational Spectroscopy, Thermal Analysis and DFT Calculations  
Zeitschrift fur Anorganische und Allgemeine Chemie  
Z. Anorg. Allg. Chem., 633 (2007), pp. 693-699, [10.1002/zaac.200600348](https://doi.org/10.1002/zaac.200600348) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [75] F.H. Allen, O. Kennard, D.G. Watson, L. Brammer, A.G. Orpen, R. Taylor  
Tables of bond lengths determined by X-ray and neutron diffraction. Part 1. Bond lengths inorganic compounds  
J. Chem. Soc. Perkin. Trans., 2S1 (1987), [10.1039/p2987000005s1](https://doi.org/10.1039/p2987000005s1) ↗  
[Google Scholar](#) ↗
- [76] Lars Wesemann, Michael Trinkaus, UlliEnglert, Jens Muller  
Silaborates with an unprecedented cluster geometry  
Organometallics, 18 (1999), pp. 4654-4659, [10.1021/om9903994](https://doi.org/10.1021/om9903994) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [77] Yuzhong Wang, Pingrong Wei YaomingXie, R. Bruce King, Henry F. Schaefer III, R. Schleyer Paul von, Gregory H. Robinson  
A stable Silicon(0) compound with a si=si double bond  
Science, 321 (2008), p. 1069, [10.1126/science.1160768](https://doi.org/10.1126/science.1160768) ↗  
[Google Scholar](#) ↗
- [78] M.J. Fink, M.J. Michalczyk, K.J. Haller, J.W.R. Michl  
X-ray crystal structures for two disilenes  
Organometallics, 3 (1984), pp. 793-800, [10.1021/om00083a025](https://doi.org/10.1021/om00083a025) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗
- [79] T. Sasamori, J.S. Han, K. Hironaka, N. Takagi, S. Nagasw, N. Tokitoh  
Synthesis and structure of stable 1,2-diaryldisilyne  
Pure Appl. Chem., 82 (2010), pp. 603-612, [10.1351/PAC-CON-09-08-02](https://doi.org/10.1351/PAC-CON-09-08-02) ↗  
[Google Scholar](#) ↗

- [80] L.V. Ya, T. Fukawa, M. Nakamoto, A. Sekiguchi, T.L. Boris, M. Karni, Y. Apeloig  
(*t*Bu<sub>2</sub>MeSi)<sub>2</sub>Sn=Sn(*SiMetBu*<sub>2</sub>)<sub>2</sub> : A Distannene with a >Sn=Sn< Double Bond That is Stable Both in the Solid State and in Solution  
J. Am. Chem. Soc., 128 (35) (2006), pp. 11643-11651, [10.1021/ja063322x](https://doi.org/10.1021/ja063322x) ↗  
[Google Scholar](#) ↗
- [81] K.W. Klinkhammer, M. Niemeyer, J. Klett  
Chem. Eur J., 5 (1999), p. 2531, [10.1002/\(SICI\)1521-3765\(19990903\)5:9%3C2531::AID-CHEM2531%3E3.0.CO;2-2](https://doi.org/10.1002/(SICI)1521-3765(19990903)5:9%3C2531::AID-CHEM2531%3E3.0.CO;2-2) ↗  
[Google Scholar](#) ↗
- [82] K. Wade  
Structural and bonding patterns in cluster chemistry  
Adv. Inorg. Chem. Radiochem., 18 (1976), pp. 1-66, [10.1016/S0065-2792\(08\)60027-8](https://doi.org/10.1016/S0065-2792(08)60027-8) ↗  
 [View PDF](#) [View article](#) [View in Scopus](#) ↗ [Google Scholar](#) ↗
- [83] K. Wade  
The structural significance of the number of skeletal bonding electron-pairs in carboranes, the higher boranes and borane anions, and various transition-metal carbonyl cluster compounds  
J. Chem. Soc. D Chem. Commun., 15 (1971), pp. 792-793, [10.1039/C29710000792](https://doi.org/10.1039/C29710000792) ↗  
[Google Scholar](#) ↗
- [84] D.M.P. Mingos  
A general theory for cluster and ring compounds of the main group and transition elements  
Nat. Phys. Sci. (Lond.), 236 (1972), pp. 99-102, [10.1038/physci236099a](https://doi.org/10.1038/physci236099a) ↗  
[Google Scholar](#) ↗
- [85] Jaromir Plesek, Tomas Jelinek, Eva Drdakova, Bohumil Stibr, Stanislav Hermanek  
A convenient preparation of 1-CB<sub>11</sub>H<sub>12</sub> and its c-amino derivatives  
Chem. Commun., 49 (1984), [10.1135/cccc19841559](https://doi.org/10.1135/cccc19841559) ↗  
[Google Scholar](#) ↗
- [86] Lauri Toom, Agnes Kutt, Ivo Leito  
Simple and scalable synthesis of the carborane anion CB<sub>11</sub>H<sub>12</sub><sup>-</sup>, dalton  
OR Trans., 48 (2019), pp. 7499-7502, [10.1039/C9DT01062A](https://doi.org/10.1039/C9DT01062A) ↗  
[View in Scopus](#) ↗ [Google Scholar](#) ↗

Cited by (0)

[View Abstract](#)

© 2025 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.



All content on this site: Copyright © 2025 Elsevier B.V., its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.

